

EQUISINGULARITY AT THE NORMALISATION

JAVIER FERNÁNDEZ DE BOBADILLA AND MARÍA PE PEREIRA

Dedicated to José María Montesinos, on the occasion of his 60th birthday

1. INTRODUCTION

We look at topological equisingularity of a holomorphic family of reduced mapping germs $f_t : (\mathbb{C}^3, O) \rightarrow \mathbb{C}$ over a contractible base T having non-isolated singularities, by means of their normalisations. The relation between surfaces in $\mathbb{C}^3$ with non-isolated singularities and normal surface singularities via the normalisation is a very convenient way of exchanging information between both categories. This has been already exploited fruitfully by T. de Jong and D. van Straten in order to study deformations of singularities [6], [7], [8].

Moreover we apply our results to the study of topological $\mathcal{A}$ -equisingularity of parametrised surfaces. The main observation is that the normalisation of a parametrised surface coincides with the parametrisation itself.

We introduce the notion of Equisingularity at the normalisation for a family f_t (see Definition 2). It turns out that in many cases, equisingularity at the normalisation characterises topological embedded equisingularity and R -equisingularity. More precisely we prove the following theorem:

Theorem A.

- If for a single $t \in T$ each of the generic transversal singularities of f_t has at least one smooth branch, then equisingularity at the normalisation is equivalent to topological R -equisingularity.
- If Lê's Conjecture holds [10], [4] (see Section 4 for the statement), then equisingularity at the normalisation is equivalent to topological R -equisingularity without any assumption on transversal singularities.

The fact that topological R -equisingularity implies equisingularity at the normalisation is not difficult and it is proved at the beginning of the paper. The proof of the converse splits in two parts. In the first part we prove the following:

Theorem B. *Equisingularity at the normalisation and equisingularity at the singular locus implies topological R -equisingularity.*

The idea of the proof is based in [3]. Here we also use the notion of *cuts*, which are real hypersurfaces in $T \times \mathbb{C}^3$ which enclose a neighbourhood of T and which imitate the situation of having a uniform radius for the Milnor fibration of f_t . Due

2000 *Mathematics Subject Classification.* Primary: 32S25, 32S50.

First author supported by Ramon y Cajal contract. Both authors supported by the Spanish project MTM2004-08080-C02-01.

to the absence of h-cobordism Theorem in the dimension we are dealing with, the techniques needed here are of a different nature. The reader may see equisingularity at the normalisation as a substitute for the h-cobordism Theorem. In the second part we search for hypothesis under which equisingularity at the normalisation implies equisingularity at the singular locus. These hypothesis are those which appeared in Theorem A.

To each family of parametrisations $\varphi_t : (\mathbb{C}^2, O) \rightarrow (\mathbb{C}^3, O)$ we can associate a family of functions $f_t : (\mathbb{C}^3, O) \rightarrow \mathbb{C}$ whose zero set coincide with the images of the family φ_t . In this situation, equisingularity at the normalisation means the constancy of the Milnor number at the origin of the inverse image by φ_t of the singular set of the image of φ_t . We prove the following theorem:

Theorem C. *Topological $\mathcal{A}$ -equisingularity of φ_t implies the constancy of the Milnor number at the origin of the inverse image by φ_t of the singular set of the image of φ_t . Conversely,*

- *if for a single $t \in T$ each of the generic transversal singularities of $\varphi_t(\mathbb{C}^2)$ has at least one smooth branch, then the constancy of the Milnor number at the origin of the inverse image by φ_t of the singular set of the image of φ_t implies topological $\mathcal{A}$ -equisingularity.*
- *If Lê's Conjecture is true, the conclusion holds in general.*

For the particular case of finitely determined mapping germs (which automatically satisfy that generic transversal singularities are of Morse type), the result was obtained also in [2]. This theorem answers a question of D. Mond in [14]. The proof of Calleja-Bedregal, Houston and Ruas uses a technique of Gaffney [5] which has the virtue of giving Whitney equisingularity as well. The finite-determinacy of germs is essential for the use of that technique. In fact we expect that in the infinite-determined case topological $\mathcal{A}$ -equisingularity does not necessarily imply Whitney equisingularity. Certainly equisingularity at the normalisation does not imply Whitney equisingularity: Briançon and Speder's family is a topologically equisingular family of normal surfaces in $\mathbb{C}^3$ (and hence equisingular at the normalisation), which is not Whitney equisingular. On the other hand our technique works under much weaker hypothesis because it uses purely topological tools instead of integration of vector fields.

Our methods also help in the study of μ -constant deformations in $\mathbb{C}^3$ since the normalisation of a μ -constant family is the identity map. Thus, Theorem B has the following corollary:

Corollary D. *If a μ -constant family of surfaces in $\mathbb{C}^3$ is topologically abstractly equisingular, then it is topologically R -equisingular.*

As a final remark, we notice that our topological trivialisations are *global* over the original base T as long as the base is contractible and compact.

2. SETTING

Let T be a contractible and compact complex manifold with a base point that is denoted by $0 \in T$. Let U be a neighbourhood of T in $T \times \mathbb{C}^3$; we denote $T \times \{0\}$ simply by T .

Throughout the article we will consider a holomorphic function

$$f : U \rightarrow \mathbb{C}$$

with $T \subset f^{-1}(0)$. As we denote with f_t the restriction of f to $\{t\} \times \mathbb{C}^3$, we will speak of a holomorphic family f_t referring to this situation. We always ask f and f_t be reduced germs at the origin for any $t \in T$.

We will call $\tau : T \times \mathbb{C}^3 \rightarrow T$ the projection onto T and denote

$$F := (\tau, f) : U \rightarrow T \times \mathbb{C}.$$

Let $(z_1, \dots, z_n)$ be a coordinate system of $\mathbb{C}^n$. Define $C := V(\partial f / \partial z_1, \dots, \partial f / \partial z_n)$. We call $X := f^{-1}(0)$ the zero set of f and $\Sigma := C \cap X$. Analogously we call $X_t := f_t^{-1}(0)$ and $\Sigma_t := \text{Sing} X_t$. Then Σ is equal to the union of the Σ_t .

In general, given any subset A contained in U we will denote $A \cap \tau^{-1}(T')$ by $A_{T'}$ for any subset T' of T . In particular, if $T' = \{t\}$ we will denote it by A_t .

For any function $g : T \times A \rightarrow B$, we will denote the function $g(t, _) : A \rightarrow B$ by g_t .

When we are working in some $\{t\} \times \mathbb{C}^3$ and the context is clear enough we will simply figure out working in $\mathbb{C}^3$.

We consider the following equisingularity notions:

Definition 1. *The family f_t is topologically equisingular if there is a homeomorphism germ*

$$(1) \quad \Phi : T \times (\mathbb{C}^3, O) \rightarrow T \times (\mathbb{C}^3, O),$$

such that we have the equality $\tau = \tau \circ \Phi$ and Φ sends X to $T \times X_0$. If in addition Φ preserves fibres of f , we say that f_t is topologically R -equisingular (the prefix R stands for right equivalence). If moreover, the restrictions $\Phi|_{T \times \mathbb{C}^3 \setminus \Sigma}$, $\Phi|_{\Sigma \setminus T}$ are smooth, we say that f_t is piecewise smoothly R -equisingular.

The family f_t is abstractly topologically equisingular if there is a homeomorphism germ

$$\Psi : (X, T) \rightarrow T \times (X_0, O)$$

such that we have the equality $\tau|_X = \tau \circ \Psi$. If in addition, Ψ sends Σ to $T \times \Sigma_0$, we say that f_t is abstractly topologically equisingular in the strong sense.

Topological R -equisingularity is stronger than topological equisingularity, which is stronger than strong abstract topological equisingularity, which is stronger than abstract topological equisingularity.

3. EQUISINGULARITY AT THE NORMALISATION

We will relate the previous equisingularity notions with the following; consider the normalisation mapping

$$n : \hat{X} \rightarrow X.$$

Consider the set $\hat{\Sigma} := n^{-1}(\Sigma)$ in which the normalisation mapping is not a local isomorphism.

In general, for any subspace A of X , the space $n^{-1}(A)$ is denoted by $\hat{A}$ and for any function $g : X \rightarrow B$, the composition $\hat{g} := g \circ n : \hat{X} \rightarrow B$ is denoted by $\hat{g}$.

Definition 2. *The family f_t is equisingular at the normalisation if there is a homeomorphism*

$$\alpha : (\hat{X}, \hat{\Sigma}, \hat{T}) \rightarrow T \times (\hat{X}_0, \hat{\Sigma}_0, n_0^{-1}(O))$$

such that we have the equality $\hat{\tau} = p_1 \circ \alpha$ where p_1 is the projection of $T \times \hat{X}_0$ to the first factor. If in addition, the restrictions $\alpha|_{\hat{X} \setminus \hat{\Sigma}}$, $\alpha|_{\hat{\Sigma} \setminus n^{-1}(O)}$ are smooth we say that f_t is piecewise smoothly equisingular at the normalisation.

(To avoid confusion we do not use the notation $\hat{O}$ or $\hat{O}_t$.)

Notation. By equisingularity at the normalisation it is immediate that, if we index the irreducible components of $\hat{\Sigma}$ as $\hat{\Sigma}^1, \dots, \hat{\Sigma}^r$, then the components of $\hat{\Sigma}_t$ can also be indexed as $\hat{\Sigma}_t^1, \dots, \hat{\Sigma}_t^r$ paralelly in the sense that $\hat{\Sigma}_t^i$ is contained in $\hat{\Sigma}^i$ for any $t \in T$.

Here are some consequences of equisingularity at the normalisation

Lemma 3. *If f_t is equisingular at the normalisation then*

- (i) *the number of irreducible components of the germ $(X, (t, O))$ is independent of t .*
- (ii) *The mapping $n_t : \hat{X}_t \rightarrow X_t$ is the topological normalisation, that is, the continuous mapping underlying it, coincides with the continuous mapping underlying the normalisation of X_t .*
- (iii) *Each connected component of $\hat{X}_t$ contains a unique point of $n_t^{-1}(O)$.*
- (iv) *For any $t \in T$, the space $(\hat{X}_t)_{\text{red}}$ is smooth outside $n_t^{-1}(O)$.*
- (v) *For any $t \in T$, the curve $(\hat{\Sigma}_t)_{\text{red}}$ is smooth outside $n_t^{-1}(O)$.*
- (vi) *If $\hat{\Sigma}_t$ is a flat, reduced family of curves then $(\hat{\Sigma}_t, n_t^{-1}(O))$ is a μ -constant family of multi-germs.*
- (vii) *If $\hat{X}_t$ is equal to $\mathbb{C}^2$ for any t , then $\hat{\Sigma}_t$ is a μ -constant family.*

Proof. The number of irreducible components of the germ $(X, (t, O))$ coincides with the number of preimages of (t, O) by the normalisation mapping, which is the cardinality of $n^{-1}(t, O)$. As $n^{-1}(t, O)$ is equal to $\alpha_t(n^{-1}(0, O))$, the first claim holds.

Since $n_t : \hat{X}_t \rightarrow X_t$ is finite and generically $1 : 1$, due to the uniqueness of the topological normalisation, it is enough to check that $\hat{X}_t$ is topologically normal at any point. By definition, the surface $\hat{X}_t$ is topologically normal if and only if $\hat{X}_t \setminus \text{Sing}(\hat{X}_t)$ is locally connected at each point of $\text{Sing}(\hat{X}_t)$. Suppose that this is not the case. The set A of points of $\text{Sing} \hat{X}_t$ where this fails, is the locally closed analytic subset of points of $\hat{X}_t$ where $\hat{X}_t$ is not locally irreducible. By equisingularity at the normalisation we can take a representative $\hat{X}$ homeomorphic through α to the product $T \times \hat{X}_t$. As any such homeomorphism transforms $T \times A$ into $\text{Sing}(\hat{X})$, we obtain that $\hat{X} \setminus \text{Sing}(\hat{X})$ is not locally connected at $\text{Sing}(\hat{X})$. This contradicts the normality of $\hat{X}$ and proves (ii).

Choosing a small enough representative of X , we can assume that (iii) is true for $t = 0$. For $t \neq 0$ it follows from the existence of α_t .

As $\hat{X}_0$ is the topological normalisation of X_0 , it has only isolated singularities. As $\hat{\Sigma}_0$ is a curve, it has only isolated singularities. Choosing a sufficiently small representative of X we may assume that $\text{Sing}(\hat{X}_0) \cup \text{Sing}(\hat{\Sigma}_0)$ is included in $n_0^{-1}(O)$. Now we use the existence of the homeomorphism α_t : by Mumford's Theorem [15] the surface $\hat{X}_t$ is smooth outside $n_0^{-1}(O)$; analogously, the curve $\hat{\Sigma}_t$ can not be singular outside $n_0^{-1}(O)$, because we would have a plane curve singularity germ topologically equivalent to a smooth one. This proves (iv) and (v).

The last assertion is a direct consequence of the sixth, which follows by the definition of equisingularity at the normalisation and the characterisation of equisingular families of flat, reduced families of curves given in [1]. $\square$

Notation. Observe that the image of each component $\hat{\Sigma}_t^i$ by n is an irreducible component of Σ_t . The restriction

$$n|_{\hat{\Sigma}_t^i} : \hat{\Sigma}_t^i \rightarrow n(\hat{\Sigma}_t^i)$$

is a finite mapping. Denote by

$$\varphi^i : \widetilde{n(\hat{\Sigma}_t^i)} \rightarrow n(\hat{\Sigma}_t^i)$$

the normalisation mapping. As $\hat{\Sigma}^i$ is topologically normal (and analytically normal outside $\hat{T}$) there is a unique finite continuous (analytic outside $\hat{T}$) mapping

$$\tilde{n}^i : \hat{\Sigma}^i \rightarrow \widetilde{n(\hat{\Sigma}^i)}$$

such that $n|_{\hat{\Sigma}_t^i} = \circ \varphi^i \circ \tilde{n}^i$. The mapping φ^i is generically $1 : 1$. We denote by d_t^i the degree of $\tilde{n}_t^i$. We define an equivalence relation R_t in the set $\{1, \dots, r\}$ defining i, j to be equivalent if we have the equality $n_t(\hat{\Sigma}_t^i) = n_t(\hat{\Sigma}_t^j)$. Then we prove the following lemma

Lemma 4. *If f_t is equisingular at the normalisation then*

- *the relation R_t is independent of t , and thus we can speak of a unique relation R .*
- *For any $t \in T$ and for any i the mapping φ_t^i is generically $1 : 1$.*
- *The degrees d_t^i are independent of t and thus, we can denote them with d^i .*
- *The fibres of $\tau|_{\Sigma} : \Sigma \rightarrow T$ are generically reduced.*

Proof. Choose a representative X small enough such that

- (a) the singular locus Σ_0 is a union of topological discs only meeting at the origin O ,
- (b) the transversal Milnor number at any point of $x \in \Sigma_0 \setminus \{O\}$ of the surface X_0 only depends on the connected component of $\Sigma_0 \setminus \{O\}$ to which x belongs,
- (c) for any irreducible component $\hat{\Sigma}_0^i$, the mapping $n|_{\hat{\Sigma}_0^i} : \hat{\Sigma}_0^i \rightarrow \Sigma_0$ is a ramified covering over its image which only ramifies at $n^{-1}(O)$.

The locus of T where $i \sim_{R_t} j$ is the set of points t such that $n(\hat{\Sigma}_t^i) = n(\hat{\Sigma}_t^j)$. Define $\Sigma_i := n(\hat{\Sigma}_i)$ for any i . Consider the restriction

$$\tau|_{\Sigma^i \cap \Sigma^j} : \Sigma^i \cap \Sigma^j \rightarrow T.$$

As Σ^i and Σ^j are irreducible, if the dimension of the fibre $(\Sigma^i \cap \Sigma^j)_t$ is not zero then Σ_t^i is equal to Σ_t^j and the fibre coincides with them. By the upper semicontinuity of the fibres of a morphism the locus where $i \sim_{R_t} j$ is closed analytic in T .

Let $A \subset n(\hat{\Sigma}^i)$ be the closed analytic subset where $n(\hat{\Sigma}^i)$ is not topologically normal. The locus of T where φ_t^i is not generically $1 : 1$ is the set where the dimension of the fibres of $\tau|_A : A \rightarrow T$ jumps to 1. This is a proper closed analytic subset.

Let $B \subset \widetilde{n(\hat{\Sigma}^i)}$ be the branching locus of the branched cover $\tilde{n}^i : \hat{\Sigma}_t^i \rightarrow \widetilde{n(\hat{\Sigma}_t^i)}$. The degree d_t^i is lower-semicontinuous in t , and the locus where it jumps down is closed analytic because it is the locus where the dimension of the fibres of the mapping $\tau|_B : B \rightarrow T$ jumps up to 1.

We have proved that there is a proper closed analytic subset $\Delta \subset T$ such that R_t and d_t^i for $1 \leq i \leq r$ are constant and φ_t^i is generically $1 : 1$ outside Δ . Moreover

R_t is finer than $R_{t'}$ and $d_t^i \geq d_{t'}^i$ if $t' \in \Delta$ and $t \in T \setminus \Delta$. In particular R_t and d_t^i are generically constant.

It is clearly enough to treat the case when T is a disc and 0 is the unique point over which the statement of the Lemma might be false.

Claim. Let $l : \mathbb{C}^3 \rightarrow \mathbb{C}$ be a generic linear function vanishing at O . Denote with L_s the hyperplane $l = s$. By genericity, L_s meets Σ_0 transversely for any $s \neq 0$ small. Then there exists a neighbourhood T' of 0 in T such that

$$(2) \quad \tau : (X \cap (T' \times L_s), \Sigma \cap (T' \times L_s)) \rightarrow T'.$$

is a μ -constant family of multigerms.

Let us prove the claim. As $n_0|_{\hat{\Sigma}_0} : \hat{\Sigma}_0 \rightarrow \Sigma_0$ is an unramified covering outside $n_0^{-1}(O)$ and Σ_0 meets L_s transversely, the mapping $n_0|_{\hat{\Sigma}_0 \setminus \{O\}}$ is transverse to L_s ; thus $n_0 : \hat{X}_0 \rightarrow \mathbb{C}^3$ is transverse to L_s at any point of $\hat{\Sigma}_0 \setminus \{O\}$. Consequently all the components of the multi-germ $(n_0^{-1}(L_s), \hat{\Sigma}_0 \cap n_0^{-1}(L_s))$ are smooth germs of curves. From now on we fix a generic s with this property.

By equisingularity at the normalisation the family $\tau : \hat{\Sigma} \rightarrow T$ is topologically equisingular. This implies easily that any fibre $\hat{\Sigma}_t$ is reduced at its generic point. As a consequence we find that the mapping

$$\hat{\tau} : \hat{\Sigma} \cap n^{-1}(T \times L_s) \rightarrow T$$

is a local diffeomorphism near $\hat{\Sigma}_0 \cap n_0^{-1}(L_s)$. Thus the mapping

$$\hat{\tau} : (n^{-1}(T \times L_s), \hat{\Sigma} \cap n^{-1}(T \times L_s)) \rightarrow T$$

is a submersion at a neighbourhood U of $\hat{\Sigma}_0 \cap n_0^{-1}(L_s)$ in $n^{-1}(T \times L_s)$. Taking local charts for the submersion we can choose U and a neighbourhood T' of 0 in T such that

$$\hat{\tau} : (U, \hat{\Sigma} \cap U) \rightarrow T'$$

is a locally trivial fibration whose fibre is diffeomorphic to the disjoint union of several punctured discs $(D, *)$. The image $(V, \Sigma \cap (T' \times L_s)) := n(U, \hat{\Sigma} \cap U)$ is a representative of the family of multi-germs of plane curve singularities

$$\tau : (X \cap (T' \times L_s), \Sigma \cap (T' \times L_s)) \rightarrow T'.$$

The mapping

$$n|_V : (U, \hat{\Sigma} \cap U) \rightarrow (V, \Sigma \cap (T' \times L_s))$$

is a simultaneous (non-embedded) resolution of singularities of the family over T' . Using this fact, we are going to prove that the Milnor number of the family of multi-germs is constant.

Observe that $\text{Sing}(V_t)$ is equal to $\Sigma_t \cap L_s$ for any $t \in T'$. The irreducible components of the multigerms $(V_t, \Sigma_t \cap L_s)$ are in a 1-1 correspondence with the discs of U_t and then its cardinality is constant for any t . We give an indexing to these components so that for any index i and any $t, t' \in T'$ the components V_t^i and $V_{t'}^i$ are images of discs lying in the same connected component of U .

Next, we prove that the intersection multiplicity in L_s of any pair of corresponding components at $\text{Sing}(V_t)$ is independent of t . If the contrary holds, there exist a pair $(V_t^1, p_t^1), (V_t^2, p_t^2)$ of components of the multigerms V_t such that the following strict inequality holds

$$I_{p_t^1}(V_t^1, V_t^2) < I_{p_0^1}(V_0^1, V_0^2)$$

for $t \neq 0$. By the conservation of the intersection number, the set $V_t^1 \cap V_t^2$ contains of a non-empty finite set of points $\{q_t^1, \dots, q_t^r\}$ different from p_t^1 and convergent to p_0^1 as t approaches 0. Each point q_t^i is a singularity of the multiterm V_t . As the singular locus of V_t is equal to $\Sigma_t \cap L_s$, and in a small enough neighbourhood W of p_0^1 in V , the only point of $\Sigma_t \cap L_s \cap W$ is p_t^1 we have got a contradiction.

Lastly, let us see that the Milnor numbers of the irreducible components are constant. Suppose that there exists a component (V_t^i, p_t) such that we have the strict inequality $\mu(V_t^i, p_t) < \mu(V_0^i, p_0)$ for $t \neq 0$. Then $\delta(V_t^i, p_t) < \delta(V_0^i, p_0)$. As V_t is a family of images of parametrisations, the sum of δ -invariants must be preserved. Therefore V_t must have singular points outside p_t which converges to p_0 as t approaches 0. So we get a contradiction and finish the proof of the claim.

Now, for any $t \in T'$ we define an equivalence relation R'_t in the finite set $\hat{\Sigma}_t \cap U$ imposing that two points are equivalents if they have the same image by n_t . The fact that the relations R'_t are independent of t follows from the equisingularity of the family (2) of plane curves. This implies easily that the fibres of $\tau|_{\Sigma} \rightarrow T$ are reduced at their generic points, that R_t is independent of t , and that φ_t^i is generically 1 : 1 for any $t \in T$. Given that the relation R_t is independent of t and that φ_t^i is generically 1 : 1 for any $t \in T$, the cardinality of the equivalence classes of R'_t only depends on the degrees d_t^i . Thus d_t^i is independent of t for any $1 \leq i \leq r$. $\square$

Lemma 5. *Let $h : X_1 \rightarrow X_2$ be a homeomorphism between reduced analytic spaces preserving the singular locus. Let $n_1 : \hat{X}_1 \rightarrow X_1$ and $n_2 : \hat{X}_2 \rightarrow X_2$ be the normalisation mappings. There is a unique homeomorphism $g : \hat{X}_1 \rightarrow \hat{X}_2$ lifting h .*

Proof. Since

$$n_i|_{\hat{X}_i \setminus \hat{\Sigma}_i} : \hat{X}_i \setminus \hat{\Sigma}_i \rightarrow X_i \setminus \Sigma_i$$

is a homeomorphism for $i = 1, 2$ We are forced to define

$$g|_{\hat{X}_1 \setminus \hat{\Sigma}_1} := n_2|_{\hat{X}_2 \setminus \hat{\Sigma}_2}^{-1} \circ h|_{X_1 \setminus \Sigma_1} \circ n_1|_{\hat{X}_1 \setminus \hat{\Sigma}_1}.$$

We only have to show that $g|_{\hat{X}_1 \setminus \hat{\Sigma}_1}$ admits a unique continuous extension to $\hat{X}_1$. Uniqueness is clear since $\hat{X}_1 \setminus \hat{\Sigma}_1$ is dense in $\hat{X}_1$. Consider $y_1 \in \hat{\Sigma}_1$. Define $x_1 := n_1(y_1)$ and $x_2 := h(x_1)$. We have to define $g(y_1)$ within the finite set $n_2^{-1}(x_2)$. The set $n_i^{-1}(x_i)$ is in a 1 : 1 correspondence with the irreducible components of the germ (X_i, x_i) for $i = 1, 2$. Thus there is a unique irreducible component (Z, x_1) of (X_1, x_1) corresponding to y_1 . We define $g(y_1) = y_2$, being y_2 the unique point of $n_2^{-1}(x_2)$ corresponding to $(h(Z), x_2)$. It is easy to check that this choice gives a continuous extension. $\square$

Proposition 6. *If f_t is abstractly topologically equisingular in the strong sense then f_t is equisingular at the normalisation.*

Proof. If Ψ is the homeomorphism which gives the abstractly topological equisingularity, by Lemma 5 there is a unique homeomorphism

$$\alpha : T \times \hat{X}_0 \rightarrow \hat{X}$$

lifting Ψ . $\square$

4. EQUISINGULARITY AT THE SINGULAR LOCUS

Definition 7. We say that f_t is equisingular at the singular locus if

- (i) the projection $\tau : \Sigma \rightarrow T$ is a (non-necessarily reduced) topologically equisingular family of curves.
- (ii) for any $t \in T$, the transversal Milnor number of f_t at any point of $\Sigma_t \setminus \{O\}$ only depends on the connected component of $\Sigma \setminus T$ in which it lies.

Observe that by the first condition of the definition, the inclusion defines a one to one correspondence between the connected components of $\Sigma_t \setminus \{O\}$ and the connected components of $\Sigma \setminus T$ for any $t \in T$.

Example (51) of [3] shows that the family Σ_t is not necessarily reduced at the origin.

In equisingularity at the singular locus we only care about the behaviour of the family of functions at the singular locus Σ of the central fibre X minus the origin. We will need also to control it at the whole critical locus C minus the origin. Thus, following [3], we make the following definition:

Definition 8. We say that f_t is equisingular at the critical locus if

- (i) it is equisingular at the singular locus and
- (ii) the spaces Σ and C coincide.

In Section 5.4 we will prove that assuming equisingularity at the normalisation, equisingularity at the singular locus implies equisingularity at the critical locus.

We recall Lê's conjecture [10] (see [4] for equivalent formulations):

Conjecture. (Lê) *The image of an injective analytic mapping germ from $(\mathbb{C}^2, O)$ to $(\mathbb{C}^3, O)$ is a surface with smooth singular locus and constant transversal Milnor number.*

Proposition 9. *The following statements hold:*

- (i) *If Lê's conjecture holds, equisingularity at the normalisation implies equisingularity at the singular locus.*
- (ii) *If for a single $t \in T$ each of the generic transversal singularities of f_t has at least one smooth branch, then the conclusion holds independently of Lê's conjecture.*

Let us state separately the following important special case of the second assertion:

Corollary 10. *If for a single $t \in T$ the generic transversal singularities of f_t are of Morse type, then equisingularity at the normalisation implies equisingularity at the singular locus.*

Proof of Proposition 9.

Step 1. We prove that the projection $\tau|_{\Sigma} : \Sigma \rightarrow T$ is a topologically equisingular family of curves, which are smooth outside of the origin. By Lemma 4 the fibres of $\tau : \Sigma \rightarrow T$ are generically reduced.

By conditions (a) – (c) of the proof of Lemma 4 and equisingularity at the normalisation we can suppose that $\hat{\Sigma}_t^i$ are discs for any t .

As $\hat{\Sigma}_t^i$ is normal we can factorise the finite mapping

$$n|_{\hat{\Sigma}_t^i} : \hat{\Sigma}_t^i \rightarrow n(\hat{\Sigma}_t^i)$$

through the normalisation $\widetilde{n(\hat{\Sigma}_t^i)} \rightarrow n(\hat{\Sigma}_t^i)$. We obtain a branched covering

$$\hat{\Sigma}_t^i \rightarrow \widetilde{n(\hat{\Sigma}_t^i)}$$

of degree d^i . Observe that, since $\widetilde{n(\hat{\Sigma}_t^i)}$ is a connected Riemann surface with boundary, its Euler characteristic must be at most 1, reaching the maximal value only when $\widetilde{n(\hat{\Sigma}_t^i)}$ is a topological disc. An Euler characteristic computation shows

$$1 = \chi(\hat{\Sigma}_t^i) = d_t^i \chi(\widetilde{n(\hat{\Sigma}_t^i)}) - R$$

where R is the total number of ramification points. We deduce that $\widetilde{n(\hat{\Sigma}_t^i)}$ is a topological disc.

Observe that Σ_t consists precisely of the union of the curves $n(\hat{\Sigma}_t^i)$ and that we know that for any $t \in T'$ all these discs meet at the origin O . For $t = 0$ we know that the origin is the unique intersection point of any of these two discs, and that any of them has no self intersections. In order to conclude the proof of Step 2 we only have to show that the same holds for any $t \in T$. Suppose the contrary: either two of the curves $n(\hat{\Sigma}_t^i)$ meet or one of them has a self intersection for a certain $t \neq 0$.

We analyse first the case in which the intersection point is not the origin: suppose that there are indexes i, j , not necessarily distinct, points $x \in \hat{\Sigma}_t^i, y \in \hat{\Sigma}_t^j$, and neighbourhoods U_x of x in $\hat{\Sigma}_{t,i}$, U_y of y in $\hat{\Sigma}_{t,j}$ such that we have the equality $n(x) = n(y)$, and the intersection $n(U_x \setminus \{x\}) \cap n(U_y \setminus \{y\})$ is empty. Consider neighbourhoods V_x of x in $\hat{X}_t$ and V_y of y in $\hat{X}_t$. The surfaces $n(V_x)$ and $n(V_y)$ in $\mathbb{C}^3$ share the common point $n(x) = n(y)$. Therefore their intersection $Z := n(V_x) \cap n(V_y)$ is a curve which is obviously contained in Σ_t . Hence $n^{-1}(Z)$ is contained in $\hat{\Sigma}$. Observe that x belongs to $n^{-1}(Z)$, but that the unique common point of $n^{-1}(Z) \cap V_x$ and $\hat{\Sigma}_t^i \cap V_x$ is the point x . Thus $\hat{\Sigma}$ is not locally irreducible at x and hence, it is singular. This contradicts Assertion (iv) of Lemma 3.

Suppose now that we have a component $\hat{\Sigma}_{t,i}$ having the origin as a self intersection point. Then there exists two distinct points $x, y \in \hat{\Sigma}_{t,i}$ such that $n(x) = n(y) = 0$. As $\hat{\Sigma}_{t,i}$ is connected, both points are in the same connected component of $\hat{X}_t$. As by equisingularity at the normalisation each of the connected components of $\hat{X}_t$ contains a unique point of $n_t^{-1}(O)$, we get a contradiction.

Step 2. We prove the second condition of equisingularity at the singular locus. As the set $n|_{\hat{\Sigma}_t^i}^{-1}(0)$ consists of a single point, we conclude that $n|_{\hat{\Sigma}_t^i}$ is the d^i -th cyclic covering of the disc onto itself totally ramified at $n|_{\hat{\Sigma}_t^i}^{-1}(0)$. Thus, n_t is locally injective at any point $y \in \hat{X}_t \setminus n_t^{-1}(0)$. By Lê's Conjecture we conclude that Σ_t (which is equal to $\cup_{i=1}^r n(\hat{\Sigma}_t^i)$) is smooth outside the origin and that for any neighborhood W of any point of $\hat{\Sigma}_t^i \setminus n_t^{-1}(0)$ in $\hat{X}_t \setminus n_t^{-1}(0)$, the image $n(W)$ is a surface possibly singular only at $n(W \cap \hat{\Sigma}_t^i)$ with constant transversal Milnor number.

In the proof of Lemma 4 we have obtained that, for any $t \in T$, the transversal Milnor number of f_t at a *generic* point of $\Sigma_t \setminus \{O\}$ only depends on the connected component of $\Sigma \setminus T$ in which it lies (this is the content of the Claim included in

the proof of the lemma). Therefore it is sufficient to show that for any $t \in T$, the transversal Milnor number of f_t at *any* point of $\Sigma_t \setminus \{O\}$ only depends on the connected component of $\Sigma_t \setminus \{O\}$ in which it lies.

Consider any point $x \in \Sigma_t \setminus \{O\}$. The number of irreducible components of the transversal singularity coincides with the number of preimages of x by n , and therefore only depends on the connected component of $\Sigma_t \setminus \{O\}$ to which x belongs. We have obtained as a consequence of Lê's Conjecture that the Milnor number of each of these irreducible components is locally constant, and hence depends only on the connected component of $\Sigma_t \setminus \{O\}$ to which x belongs. It only remains to be shown that the intersection multiplicities between the different branches of the transversal singularities are locally constant in $\Sigma_t \setminus \{O\}$. Such intersection multiplicities are upper semicontinuous.

Suppose that there is a special point $x \in \Sigma_t \setminus \{O\}$ in which the intersection multiplicity between two branches jumps. More precisely, given a linear function $l : \mathbb{C}^3 \rightarrow \mathbb{C}$ meeting Σ_t transversely at x , the branches of the plane curve singularity given by the section defined by l at a point $x' \in \Sigma_t$ close to x are in bijective correspondance with the set of preimages of x' by n_t . There are two points $y_1, y_2 \in n_t^{-1}(x)$ such that the intersection multiplicity of the two associated branches of the hyperplane section is strictly bigger than the intersection multiplicity of the corresponding branches at neighbouring hyperplane sections. For $i = 1, 2$ we choose a small neighbourhood V_{y_i} of y_i in $\hat{X}_t$, with a biholomorphism $\varphi_i : V_{y_i} \rightarrow D \times D$ (denoting D the disc) such that the intersection $\varphi_i(\hat{\Sigma}_t \cap V_{y_i})$ is equal to $\{0\} \times D$ and that $\varphi_i^{-1}(D \times \{a\})$ for $a \in D$ are the inverse images by n of the hyperplane sections by L . By the conservation of the intersection multiplicity, for a certain $a \in D$, the two branches $n(\varphi_i^{-1}(D \times \{a\}))$ for $i = 1, 2$ must intersect outside Σ . Their intersection points are singular points of X_t , and therefore must belong to Σ . We conclude that the intersection multiplicity is locally constant. This concludes Step 3 and the proof of Assertion (i).

We prove Assertion (ii). In Step 1 we have proved that for any $t \in T$, the transversal Milnor number of f_t at a *generic* point of $\Sigma_t \setminus \{O\}$ only depends on the connected component of $\Sigma \setminus T$ in which it lies. Using this and the fact that for a single t each of the generic transversal singularities of f_t at $\Sigma_t \setminus \{O\}$ has at least one smooth branch, we deduce that for *any* $t \in T$ every generic transversal singularity of f_t at $\Sigma_t \setminus \{O\}$ has at least one smooth branch.

We observe that we have used Lê's conjecture only at Step 2. Using the existence of smooth branches, we can replace the use of Lê's conjecture by Mumford's Theorem (see [15]) which states that the unique normal surface germ having a link with trivial fundamental group is the smooth germ, and prove that Σ_t is smooth outside of the origin. Then we can use the fact that Lê's conjecture holds if the image contains a smooth branch [4] and proceed as in Step 2. $\square$

5. TOPOLOGICAL EQUISINGULARITY

The purpose of this section is to prove the following theorem:

Theorem 11. *Let f_t be equisingular at the singular locus and equisingular at the normalisation, then it is R -topologically equisingular.*

To prove this, we follow the idea of [3] and base our proof on constructing a family of hypersurfaces in $T \times \mathbb{C}^3$, that we will call *cuts*, which enclose neighbourhoods of T where the restriction of τ is a trivial fibration and where, near Σ , we recognise a μ -constant family of isolated singularities. We use the global version of the Theorem of Lê-Ramanujan in [3] to trivialise this μ -constant family and use the equisingularity at the normalisation to trivialise outside that neighbourhood of Σ .

For the rest of the section the family f_t will be equisingular at the singular locus and equisingular at the normalisation.

5.1. Cuts.

Notation. Let $E \rightarrow N$ be an arbitrary vector bundle with a metric ρ in its fibres. Given any function $\nu : N \rightarrow \mathbb{R}_+$, for any $N' \subset N$ we call

$$E(N', \nu) := \{(x, v) \in E_{N'} : \rho_x(0, v) < \nu(x)\}.$$

For the bundle τ and its restrictions we use the notation $B(T, \nu)$.

Assuming equisingularity at the singular locus, we can fix a complex normal bundle $\pi : E \rightarrow \Sigma \setminus T$ in $T \times \mathbb{C}^3$ and a positive function $\mu : \Sigma \setminus T \rightarrow \mathbb{R}_+$, such that $E(\Sigma \setminus T, \mu)$ can be embedded in $T \times \mathbb{C}^3$. The restriction

$$\pi|_{E(\Sigma \setminus T, \mu)} : E(\Sigma \setminus T, \mu) \rightarrow \Sigma \setminus T$$

is a fibration with fibre a ball in $\mathbb{C}^2$. Observe that the function f restricted to this bundle is a μ -constant family of curves.

Definition 12. A topological hypersurface H in $T \times \mathbb{C}^3$ is a cut for Σ of amplitude δ if

- (i) the intersection $H \cap E(\Sigma \setminus T, \mu)$ coincides with $E(H \cap \Sigma, \mu)$,
- (ii) the hypersurface H is topologically transversal to $F^{-1}(t, s)$ outside Σ , for any $(t, s) \in T \times D_\delta$,
- (iii) the hypersurface H separates $F^{-1}(T \times D_\delta)$ in several connected components, one of them is a bounded neighbourhood of T ; we call it $\text{Int}H(\delta)$. (We will not specify the amplitude if it is clear from the context). Besides, $\overline{\text{Int}H(\delta)}$ is a topological manifold with boundary such that

$$\tau|_{\overline{\text{Int}H(\delta)}} : \overline{\text{Int}H(\delta)} \rightarrow T$$

is a topological submersion.

- (iv) for any $t \in T$, $(\text{Int}H(\delta) \cap \Sigma)_t$ is homeomorphic to a cone over $H \cap \Sigma_t$.

Abusing notation slightly, we consider the subspace $\hat{E}(\mu) := n^{-1}(E(\Sigma \setminus T, \mu))$ which inherits the unique natural bundle structure

$$(3) \quad \hat{\pi} : \hat{E}(\mu) \rightarrow \hat{\Sigma} \setminus \hat{T}$$

compatible with the one in $E(\Sigma \setminus T, \mu)$.

Definition 13. A topological real hypersurface $\hat{H}$ in $\hat{X}$ is a cut for $\hat{\Sigma}$ in $\hat{X}$ if

- (i) The restriction $\hat{\tau}|_{\hat{H}}$ is a proper topological submersion.
- (ii) For any $t \in T$ the intersection $(\hat{H} \cap \hat{\Sigma})_t$ is a smooth real curve saturated by n , that is $n^{-1}(n((\hat{H} \cap \hat{\Sigma})_t))$ coincides with $(\hat{H} \cap \hat{\Sigma})_t$.
- (iii) For any $t \in T$ the smooth curve $n(\hat{H} \cap \hat{\Sigma})_t$ separates Σ_t in several connected components. The only component containing the origin is a cone over the curve $n(\hat{H} \cap \hat{\Sigma})_t$.

- (iv) The intersection $\hat{H} \cap \hat{E}(\mu)$ coincides with $\hat{\pi}^{-1}(\hat{H} \cap \hat{\Sigma})$.
- (v) The cut $\hat{H}$ separates $\hat{X}$ in several connected components, the union of those meeting $\hat{T}$ is a neighbourhood of $\hat{T}$ that we call $\text{Int}\hat{H}$. Any connected component of $(\text{Int}\hat{H})_t$ is a cone over the only point of $n_t^{-1}(0)$ that it contains.

Note. The notation of $\hat{H}$ is due to the fact that, as the reader will see in Section 5.5, a cut $\hat{H}$ can be the preimage by the normalisation n of a cut H .

In order to simplify the exposition we do the following general definition:

Definition 14. Let A be a subspace of U (resp. $\hat{X}$) that contains a subset T' of T (resp. $\hat{T}' \subset \hat{T}$). Let M a subspace of A which does not meet T' (resp. $\hat{T}'$). We can define

$$(\text{Int}_A M)_{T'}$$

as the union of the connected components of $(A \setminus M)_{T'}$ meeting T' (resp. with $\hat{T}'$). $(\text{Int}_A M)_{T'}(\delta_i)$ will stand for $(\text{Int}_{A \cap f^{-1}(D_{\delta_i})} M)_{T'}$.

When the context makes clear who A and T' are, we will not specify them. Observe that this definition is coherent with the definition of $\text{Int}H$ and $\text{Int}\hat{H}$ given above for certain cuts H and $\hat{H}$: $\text{Int}H = (\text{Int}_U H)_T$, $\text{Int}\hat{H} = (\text{Int}_{\hat{X}} \hat{H})_{\hat{T}}$. Also, as we will do in the Section 5.3, taking a hypersurface M in $\Sigma \setminus T$ we can consider $\text{Int}M = (\text{Int}_{\Sigma} M)_T$.

5.2. A glueing technique. In this subsection we explain a technical tool that we use throughout the rest of the paper. In our proofs we will have to construct piecewise several homeomorphisms. We will be working on spaces fibred over the contractible space of parameters T and most of our constructions in each of the pieces will be trivial over the parameter $0 \in T$. We will show in this subsection how to use the retraction of T onto 0 to make them compatible.

Let $(\Lambda, 0)$ be a contractible and compact topological space. Let A be a topological space. Let $f_i : A \rightarrow B_i$ be continuous mappings to topological spaces B_i , for $i \in \{1, \dots, m_1\}$ and C_i, D_j be subspaces of A , for $i \in \{1, \dots, m_2\}$, $j \in \{1, \dots, m_3\}$.

Let

$$\tilde{h} : \Lambda \times A \rightarrow \Lambda \times A$$

be a self-homeomorphism of the form $\tilde{h} := (Id|_{\Lambda}, h_{\lambda})$ with $h_{\lambda} : A \rightarrow A$ a family of homeomorphisms depending continuously in $\lambda \in \Lambda$. Suppose furthermore that this family of homeomorphisms satisfies the following conditions:

- (a) the homeomorphism $\tilde{h}_0$ is the identity,
- (b) the fibres of each of the mappings $f_i : A \rightarrow B_i$ are preserved by $\tilde{h}_{\lambda}$ for all $\lambda \in \Lambda$,
($f_i \circ \tilde{h}_{\lambda} = f_i$),
- (c) the subspaces C_i are left invariant by $\tilde{h}_{\lambda}$ for all $\lambda \in \Lambda$,
- (d) the subspaces D_j are fixed by $\tilde{h}_{\lambda}$ for all $\lambda \in \Lambda$.

We associate any continuous mapping

$$g : (\Lambda, 0) \rightarrow (\Lambda, 0)$$

with a homeomorphism

$$\tilde{g} : \Lambda \times A \rightarrow \Lambda \times A$$

defined by $\tilde{g}(\lambda, x) := (\lambda, \tilde{h}_{g(\lambda)}(x))$; the homeomorphisms $\tilde{g}_{\lambda}$ obviously satisfy conditions (a)-(d).

Lemma 15. *Let $\tilde{h} : \Lambda \times A \rightarrow \Lambda \times A$ be a homeomorphism as above. There exists an isotopy*

$$\phi : \Lambda \times A \times I \rightarrow \Lambda \times A \times I$$

between $\phi_0 = \tilde{h}$ and $\phi_1 = Id_{\Lambda \times A}$ of the form

$$\phi(\lambda, x, t) = (\lambda, \phi_{(\lambda, t)}(x), t)$$

for certain family of homeomorphisms $\phi_{(\lambda, t)} : A \rightarrow A$ which satisfy (a)-(d).

Proof. As Λ is contractible, there exists a strong deformation retract of Λ to the point 0

$$\xi : \Lambda \times I \rightarrow \Lambda$$

which satisfies $\xi(\lambda, 0) = \lambda$, $\xi(\lambda, 1) = 0$ for all $\lambda \in \Lambda$ and $\xi(0, t) = 0$ for all $t \in [0, 1]$.

We define ϕ_t as the homeomorphism $\tilde{\xi}_t$ associated to ξ_t and $\phi := (\phi_t, Id|_{[0,1]})$.

It is clear that the mapping ϕ is bijective and continuous. Its inverse is continuous because the family of homeomorphisms $\tilde{h}_\lambda^{-1}$ depends continuously in Λ (for being $\tilde{h}^{-1} : \Lambda \times A \rightarrow \Lambda \times A$ continuous.). It is obvious that ϕ satisfies the rest of the requirements. $\square$

Let E be another topological space and $\beta : E \rightarrow \Lambda$ a continuous mapping. Consider continuous mappings $f_i : E \rightarrow B_i$ to topological spaces B_i for $1 \leq i \leq m_1$ and subspaces $C_i \subset E$ for $1 \leq i \leq m_2$.

Definition 16. *A trivialisation of $\beta : E \rightarrow \Lambda$ relative to $f_i : E \rightarrow B_i$ for $1 \leq i \leq m_1$ and $C_i \subset E$ for $1 \leq i \leq m_2$ is a homeomorphism*

$$\xi : E \rightarrow \Lambda \times E_0$$

with $E_0 = \beta^{-1}(0)$ such that

- *we have the equality $\beta = p_1 \circ \xi$, where p_i denotes the projection of $\Lambda \times E_0$ to the i -th factor,*
- *it satisfies $\xi|_{E_0} = (0, Id|_{E_0})$,*
- *we have the equality $f_i = f_i|_{E_0} \circ p_2 \circ \xi$ for $1 \leq i \leq m_1$,*
- *we have the equality $\xi(C_i) = \Lambda \times (E_0 \cap C_i)$ for $1 \leq i \leq m_2$.*

Example 17. *The definition of equisingularity at the normalisation can be stated saying that $\hat{\tau} := \tau \circ n : \hat{X} \rightarrow T$ admits a trivialisation $\alpha : \hat{X} \rightarrow T \times \hat{X}_0$ relative to $\hat{\Sigma}$ and $\hat{T}$.*

Example 18. *If $\beta : E \rightarrow \Lambda$ is a proper differentiable submersion, then the Ehresman Fibration Theorem tells that there exist a trivialisation of β .*

Example 19. *Analogously, if $\beta : E \rightarrow \Lambda$ is a proper topological submersion, then there exist also a trivialisation of β (see [9] page 59).*

Assume that E is the union of two closed subspaces E_1 and E_2 . Suppose that for $i = 1, 2$ we have trivialisations of $\beta|_{E_i}$

$$\xi_i : E_i \rightarrow \Lambda \times (E_i)_0$$

relative to $f_i|_{E_i}$ for $1 \leq i \leq m_1$ and to the subsets $C_j \cap E_i$ for $1 \leq j \leq m_2$ and $E_1 \cap E_2$. We want to modify ξ_1 and ξ_2 so that they can be plumbed together to give a trivialisation of E relative to f_i for $1 \leq i \leq m_1$ and C_i for $1 \leq i \leq m_2$.

Definition 20. Let A be a subspace of E . We say that a neighbourhood $C(A)$ of E is a collar for A in E relative to f_i for $1 \leq i \leq m_1$ and to the subsets C_i for $1 \leq i \leq m_2$, if there is a homeomorphism

$$\theta : A \times [0, 1] \rightarrow C(A)$$

such that

- its restriction θ_0 is the inclusion of A in $C(A)$,
- it satisfies $f_i \circ \theta = f_i \circ p_1$ for $1 \leq i \leq m_1$, with $p_1 : A \times [0, 1] \rightarrow A$ the projection onto the first factor,
- we have the equality $\theta((C_i \cap A) \times [0, 1]) = C_i \cap C(A)$.

Lemma 21. If $A := (E_1 \cap E_2)_0$ has a collar $C(A)$ in $(E_1)_0$ relative to $f_i|_{(E_1)_0}$ for $1 \leq i \leq m_1$ and relative to the subsets $C_i \cap (E_1)_0$ for $1 \leq i \leq m_2$, then the trivialisations ξ_1 and ξ_2 can be plumbed in order to obtain a trivialisation of $\beta : E \rightarrow \Lambda$,

$$\xi : E \rightarrow \Lambda \times E_0,$$

relative to f_i for $1 \leq i \leq m_1$ and C_i for $1 \leq i \leq m_2$ and such that $\xi|_{E_2} = \xi_2$ and $\xi|_{E_1 \setminus \xi_1^{-1}(\Lambda \times C(A))} = \xi_1$.

Proof. The homeomorphism

$$h := \xi_2|_{E_1 \cap E_2} \circ \xi_1^{-1}|_{\Lambda \times (E_1 \cap E_2)_0} : \Lambda \times (E_1 \cap E_2)_0 \rightarrow \Lambda \times (E_1 \cap E_2)_0$$

satisfies the hypothesis of Lemma 15 so there exists an isotopy

$$\phi : \Lambda \times (E_1 \cap E_2)_0 \times [0, 1] \rightarrow \Lambda \times (E_1 \cap E_2)_0 \times [0, 1]$$

with $\phi_0 = h$ and $\phi_1 = Id_{\Lambda \times A}$.

We call θ the homeomorphism which gives $C(A)$ the structure of collar of A in $(E_1)_0$ relative to f_i for $1 \leq i \leq m_1$ and C_i for $1 \leq i \leq m_2$.

We can construct $\xi : E \rightarrow \Lambda \times E_0$ piecewise as follows

$$\begin{aligned} \xi|_{E_2} &:= \xi_2, \\ \xi|_{E_1 \setminus \xi_1^{-1}(\Lambda \times C(A))} &:= \xi_1, \\ \xi|_{\xi_1^{-1}(\Lambda \times C(A))} &:= (Id_\Lambda, \theta) \circ \phi \circ (Id_\Lambda, \theta^{-1}) \circ \xi_1. \end{aligned}$$

□

5.3. Existence of cuts for $\hat{\Sigma}$. In proving the existence of cuts for $\hat{\Sigma}$ we will use several auxiliary hypersurfaces in Σ that we construct thanks to the following lemma:

Lemma 22. Suppose that f_t is equisingular at the singular locus. Given any positive function $\eta : T \rightarrow \mathbb{R}_+$. Choose $\epsilon_* > 0$ such that $\mathbb{S}_\epsilon$ is transverse to Σ_0 for any $\epsilon < \epsilon_*$. There exists $0 < \epsilon < \epsilon_*$ and a smooth hypersurface M of $\Sigma \setminus T$ such that

- (i) the restriction $\tau|_M$ is a submersion,
- (ii) it separates Σ in several connected components, the only component $IntM$ containing T is contained in $B(T, \eta)$,
- (iii) for any $t \in T$ the fibre M_t is a disjoint union of circles, one located in each of the irreducible components of Σ_t ,
- (iv) for any $t \in T$ the fibre $IntM_t$ is homeomorphic to a cone over M_t ,
- (v) the fibre M_0 is equal to $\mathbb{S}_\epsilon \cap \Sigma_0$.

Proof. The proof is an easy application (actually a simplification) of the technique of construction of cuts developed in [3], Section 2. □

In this subsection we prove the following proposition:

Proposition 23. *Let f_t be equisingular at the singular locus and equisingular at the normalisation. Given any positive function $\eta : T \rightarrow \mathbb{R}_+$ there exist a cut $\hat{H}$ for $\hat{\Sigma}$ such that $n(\text{Int}\hat{H}) \subset B(T, \eta)$.*

Proof.

We construct first a cut for $\hat{\Sigma}$ over 0 and then extend it over T using the homeomorphism α^{-1} .

Choose $0 < \epsilon_* < \eta(0)$ such that the sphere $\mathbb{S}_\epsilon$ is transverse (in the stratified sense) to X_0 for any $\epsilon \leq \epsilon_*$.

We can take $0 < \epsilon_0 < \epsilon_*$ such that $\alpha^{-1}(T \times n_0^{-1}(B_{\epsilon_0}))$ is contained in $n^{-1}(B(T, \eta))$.

Using the technique of [3], Section 3, we modify $\mathbb{S}_{\epsilon_0}$ to get a neighbourhood W of the origin in $\mathbb{C}^3$ such that

- the hypersurface $\alpha^{-1}(T \times n_0^{-1}(\partial W))$ is contained in $n^{-1}(B(T, \eta))$,
- we have the equality $\partial W \cap \Sigma_0 = \mathbb{S}_{\epsilon_0} \cap \Sigma_0$,
- the boundary ∂W coincides with the normal bundle E in a neighbourhood of $\Sigma_0 \cap \partial W$ in ∂W . Precisely, we have the equality

$$\partial W \cap E(\Sigma \setminus T, \mu)_0 = E(\partial W \cap \Sigma_0, \mu).$$

The arguments needed here are much simpler than those in [3] since we have to ensure neither smoothness of ∂W nor any cobordism condition.

The topological hypersurface

$$(4) \quad L := \alpha^{-1}(T \times n_0^{-1}(\partial W))$$

is a candidate for being a cut. It satisfies the properties (i) and (v) of Definition 13 and, over the parameter 0, also the other ones. The rest of the proof consists in modifying α so that the cut L , defined as in (4), satisfies all the required conditions over any $t \in T$.

For the rest of the proof we will need three auxiliary hypersurfaces of $\Sigma \setminus T$, that we will call M_+ , M_* and M_- , easily constructed applying Lemma 22, satisfying the following conditions:

- We have the inclusions

$$\text{Int}_\Sigma M_- \subset \text{Int}_\Sigma M_* \subset \text{Int}_\Sigma M_+$$

$$\text{Int}_{\hat{\Sigma}} \hat{M}_- \subset \text{Int}_{\hat{\Sigma}} L \subset \text{Int}_{\hat{\Sigma}} \hat{M}_+$$

- The restriction $(M_*)_0$ coincides with $\mathbb{S}_{\epsilon_0} \cap \Sigma_0$,

Observe that $\hat{M}_-$, $\hat{M}_*$ and $\hat{M}_+$ are smooth topological hypersurfaces of $\hat{\Sigma}$ and that we have the equalities $\text{Int}_{\hat{\Sigma}} \hat{M}_- = n^{-1}(\text{Int}_\Sigma M_-)$, $\text{Int}_{\hat{\Sigma}} \hat{M}_* = n^{-1}(\text{Int}_\Sigma M_*)$ and $\text{Int}_{\hat{\Sigma}} \hat{M}_+ = n^{-1}(\text{Int}_\Sigma M_+)$. Observe also that the hypersurface $\hat{M}_*$ satisfies conditions (i) – (iii) of Definition 13.

For the rest of the proof we will assume that $\hat{\Sigma}$ is irreducible. The general case follows similarly.

Step 1. We modify α so that L , defined as in (4), satisfies conditions (i) – (iii) and (v). Define $P := \overline{\text{Int}\hat{M}_+} \setminus \text{Int}\hat{M}_-$. The restriction $\hat{\tau}|_P : P \rightarrow T$ is a trivial fibration with fibre a cylinder (in the general case, as many cylinders as irreducible components $\hat{\Sigma}$ has). This is true by condition (iv) of Lemma 22 and the fact that the restriction $n|_{\Sigma \setminus \hat{T}}$ is a covering projection.

Consider $Q := n^{-1}(\pi|_{E(\Sigma \setminus T, \mu)}^{-1}(\overline{\text{Int}M_+ \setminus \text{Int}M_-}))$ and the inherited bundle structure $\hat{\pi}|_Q := Q \rightarrow P$. The fibre is a disc $D \subset \mathbb{C}$ and the bundle is trivial because it is a orientable fibration over P , which has the homotopy type of $\mathbb{S}^1$. Since the bundle is trivial we identify Q with $P \times D$, noting that the product structure is compatible with the fibration $\hat{\tau}$.

We show that there exists a homeomorphism

$$\Theta : Q \rightarrow Q$$

such that it preserves the fibres of $\hat{\tau}$, it is the identity over Q_0 , it preserves P , it carries $L \cap P$ to $\hat{M}_*$ and it leaves ∂Q fixed.

Suppose that this homeomorphism exists. We extend it with the identity outside Q obtaining a homeomorphism

$$\bar{\Theta} : \hat{X} \rightarrow \hat{X}.$$

Since $\bar{\Theta}$ transforms $L \cap P$ in $\hat{M}_*$, the topological hypersurface $\bar{\Theta}(L)$ satisfies Properties (i) – (iii) and (v) of Definition 13. Redefining $\alpha := \bar{\Theta} \circ \alpha$ we can assume that L , defined as in (4), satisfies Properties (i) – (iii) and (v).

To construct Θ , we construct first an isotopy

$$\theta : P \times [0, 1] \rightarrow P \times [0, 1]$$

such that it preserves fibres of $\hat{\tau}$, the restriction θ_1 is equal to Id_P , the restriction θ_0 carries $L \cap P$ to $\hat{M}_*$ and it leaves $\partial P = \hat{M}_+ \cup \hat{M}_-$ fixed.

Then Θ can be easily constructed using θ as follows. For any $(p, z) \in P \times D = Q$ we define

$$\Theta(p, z) := (\theta|_z(p), z).$$


Let us construct θ . First of all, consider a trivialisation of $\hat{\tau}|_P : P \rightarrow T$

$$\beta : P \rightarrow T \times P_0$$

and a product structure for P_0 ,

$$\gamma : P_0 \rightarrow \mathbb{S}^1 \times [0, 1]$$

that carries $(\hat{M}_+)_0$ to $\mathbb{S}^1 \times \{0\}$, $(L \cap P)_0$ (which is equal to $(\hat{M}_*)_0$) to $\mathbb{S}^1 \times \{\frac{1}{2}\}$ and $(\hat{M}_-)_0$ to $\mathbb{S}^1 \times \{1\}$.



By construction, the hypersurface L separates P in two connected components, $P_{1,1}$ and $P_{1,2}$. Observe that both $\partial P_{1,1}$ and $\partial P_{1,2}$ have two connected components, $\hat{M}_+$ and L for the first one, and L and $\hat{M}_-$ for the second one. The restrictions

$\hat{\tau}|_{P_{1,1}}$ and $\hat{\tau}|_{P_{1,2}}$ are trivial fibrations with fibre a cylinder so that we can take trivialisations $\sigma_{1,1} : P_{1,1} \rightarrow T \times \mathbb{S}^1 \times [0, \frac{1}{2}]$ which carries M_+ to $T \times \mathbb{S}^1 \times \{0\}$ and L to $T \times \mathbb{S}^1 \times \{\frac{1}{2}\}$, $\sigma_{1,2} : P_{1,2} \rightarrow T \times \mathbb{S}^1 \times [\frac{1}{2}, 1]$ which carries L to $T \times \mathbb{S}^1 \times \{\frac{1}{2}\}$ and $\hat{M}_-$ to $T \times \mathbb{S}^1 \times \{1\}$.

We can glue them along $T \times \mathbb{S}^1 \times \{\frac{1}{2}\}$ with the mapping $\sigma_{1,2} \circ \sigma_{1,1}^{-1}$ to obtain a homeomorphism

$$\sigma_1 : P \rightarrow T \times \mathbb{S}^1 \times [0, 1] = (T \times \mathbb{S}^1 \times [0, \frac{1}{2}]) \bigcup_{\sigma_{1,2} \circ \sigma_{1,1}^{-1}} (T \times \mathbb{S}^1 \times [\frac{1}{2}, 1])$$

which carries $\hat{M}_+$ to $T \times \mathbb{S}^1 \times \{0\}$, $L \cap P$ to $T \times \mathbb{S}^1 \times \{\frac{1}{2}\}$ and $\hat{M}_-$ to $T \times \mathbb{S}^1 \times \{1\}$.

The hypersurface $\hat{M}_*$ also separates P in two connected components, $P_{2,1}$ and $P_{2,2}$. In this case, their frontiers have components $\hat{M}_+$ and $\hat{M}_*$, and $\hat{M}_*$ and $\hat{M}_-$ respectively. Analogously to the construction of σ_1 made for L we can obtain a homeomorphism

$$\sigma_2 : P \rightarrow T \times \mathbb{S}^1 \times [0, 1]$$

which carries $\hat{M}_+$ to $T \times \mathbb{S}^1 \times \{0\}$, $\hat{M}_*$ to $T \times \mathbb{S}^1 \times \{\frac{1}{2}\}$ and $\hat{M}_-$ to $T \times \mathbb{S}^1 \times \{1\}$.

Consider the homeomorphism

$$h := \beta \circ \sigma_2^{-1} \circ \sigma_1 \circ \beta^{-1} : T \times P_0 \rightarrow T \times P_0$$

which satisfies that the restriction h_0 is the identity, it carries $\beta(L \cap P)$ to $\beta(\hat{M}_*)$ and it preserves the fibres of the projection $p_1 : T \times P_0 \rightarrow T$.

Using Lemma 21 we can modify σ_2 in collars of M_+ , M_- in P such that h also leaves $T \times (M_+)_0$ and $T \times (M_-)_0$ fixed.

Applying now Lemma 15, there is an isotopy

$$\phi : T \times P_0 \times [0, 1] \rightarrow T \times P_0 \times [0, 1]$$

which satisfies that $\phi_0 = h$, $\phi_1 = Id_{T \times P_0}$ and that ϕ_s preserves the fibres of the projection $p_1 : T \times P_0 \rightarrow T$ and leaves $T \times (M_+)_0$ and $T \times (M_-)_0$ fixed for any $s \in [0, 1]$.

We finish defining θ as the composition

$$(\sigma_2^{-1}, Id_{[0,1]}) \circ (Id_T, \gamma, Id_{[0,1]}) \circ \phi \circ (Id_T, \gamma^{-1}, Id_{[0,1]}) \circ (\sigma_1, Id_{[0,1]}).$$

Step 2. We modify again α in order to make L , defined as in (4), satisfy also condition (iv) of Definition 13.


Take two tubular neighbourhoods $G_- \subset G_+$ of $L \cap \hat{\Sigma}_0$ in $\hat{\Sigma}_0$. Define $(K_-)_0 := \hat{\pi}^{-1}(G_-) \cap \hat{E}(\eta_-)$ and $(K_+)_0 := \hat{\pi}^{-1}(G_+) \cap \hat{E}(\eta_+)$ for positive functions $\eta_-, \eta_+ : \Sigma \setminus T \rightarrow \mathbb{R}^+$ such that $\eta_- < \eta_+ < \mu$. Define

$$K_+ := \alpha^{-1}(T \times (K_+)_0),$$

$$K_- := \hat{\pi}^{-1}(\alpha^{-1}(T \times G_-)) \cap \hat{E}(\eta_-),$$

$$L_* := \hat{\pi}^{-1}(L \cap \hat{\Sigma}) \cap \hat{E}(\eta_-).$$

Notice that $(L_*)_0$ coincides with $L_0 \cap \hat{E}(\eta_-)$. If η_- is sufficiently small then we have the inclusion $K_- \subset \dot{K}_+$.



By construction of K_+ we have a trivialisation

$$\alpha|_{\hat{X} \setminus K_+} : \hat{X} \setminus K_+ \rightarrow T \times (\hat{X} \setminus K_+)_0.$$

of $\tau|_{\hat{X} \setminus K_+}$, and, as $\tau|_{K_+ \setminus K_-}$ and $\tau|_{K_-}$ are proper topological submersions, they admit trivialisations

$$\kappa_1 : (K_+ \setminus K_-) \rightarrow T \times ((K_+ \setminus K_-)_0)$$

$$\kappa_2 : (K_-, L_*) \rightarrow T \times ((K_-)_0, (L \cap K_-)_0).$$

Taking collars of ∂K_+ and ∂K_- in $K_+ \setminus K_-$, we can apply Lemma 21 and obtain a trivialisation

$$\kappa : \hat{X} \rightarrow T \times \hat{X}_0$$

such that $\kappa|_{\hat{X} \setminus K_+} = \alpha|_{\hat{X} \setminus K_+}$ and $\kappa|_{K_-} = \kappa_2$.

Redefining $\alpha := \kappa$ and defining $\hat{H}$ as in (4), we have the cut required. $\square$

5.4. Equisingularity at the critical locus.

Proposition 24. *If f_t is equisingular at the singular locus and equisingular at the normalisation, then it is equisingular at the critical locus.*

Proof. We have to prove that the polar variety $\Gamma := \overline{C \setminus \Sigma}$ is empty. It is clearly sufficient to assume that T is a disc. By the second property of equisingularity at the singular locus, Thus Γ is a curve. If Γ does not meet T we obtain the conclusion shrinking the representative U . Suppose that Γ meets T at a point t_0 . We may choose the base point 0 to be equal to t_0 . Denote by μ_0 the intersection multiplicity $I_{(0,0)}(\Gamma, \{0\} \times \mathbb{C}^3)$.

Choose radii ϵ, δ for the Milnor fibration of the function f_0 (in particular, $B_\epsilon \cap f_0^{-1}(D_\delta)$ is contractible). The arguments given in the proof of Lemma 26 of [3] show that equisingularity at the singular locus implies the existence of a small neighbourhood T' of 0 in T such that $F^{-1}(t, s)$ is transversal to $\mathbb{S}_\epsilon$ (in the stratified sense if $s = 0$) for any $(t, s) \in T' \times D_\delta$. If T' is small enough, for any $t \in T' \setminus \{0\}$, the function f_t has isolated critical points outside $f_t^{-1}(0)$ whose Milnor numbers add up μ_0 . Under these conditions it follows by standard techniques that $X_t \cap B_\epsilon$ has the homotopy type of a bouquet of μ_0 spheres of dimension 2.

Using Proposition 23 we construct a cut $\hat{H}$ for $\hat{\Sigma}$ such that $\hat{H}_0$ is equal to $n_0^{-1}(\mathbb{S}_{\epsilon/2})$. If T' is small enough we have that $n(\text{Int} \hat{H})_t$ is contained in $\hat{B}_\epsilon$ for any $t \in T'$. We claim that the inclusion

$$n(\hat{H})_t \hookrightarrow \overline{X_t \cap B_\epsilon \setminus n(\text{Int} \hat{H})_t}$$

is a homotopy equivalence.

If the claim is true, then $B_\epsilon \cap X_t$ has the same homotopy type as $X_t \cap n(\text{Int}\hat{H})_t$. For any $t \in T$, the pair $\text{Int}\hat{H} \cap (\hat{X}_t, \hat{\Sigma}_t)$ is a disjoint union of cones, whose vertices are the elements of $n_t^{-1}(t)$. From the topological viewpoint the mapping

$$n_t : \hat{X}_t \cap \text{Int}\hat{H} \rightarrow n_t(\hat{X}_t \cap \text{Int}\hat{H})$$

is a quotienting of $\hat{X}_t \cap \text{Int}\hat{H}$ which can be factorised as the composition of two identifications

$$\hat{X}_t \cap \text{Int}\hat{H} \xrightarrow{a} Z \xrightarrow{b} n_t(\hat{X}_t \cap \text{Int}\hat{H})$$

which have the following properties.

- The restriction $a|_{\hat{X}_t \cap \text{Int}\hat{H} \setminus \hat{\Sigma}_t}$ is injective. Let $\{\hat{\Sigma}_t^i\}_{i=1}^d$ be the irreducible components of $\hat{\Sigma}_t$. For any $i \leq d$ there is a cyclic action on $\hat{\Sigma}_t^i$ of $\hat{\Sigma}_t$, being the vertex the only point with non-trivial isotropy group such that the fibres of a over any point of $a(\hat{\Sigma}_t^i)$ coincide with the orbits of the action.
- The restriction $b|_{Z \setminus a(\hat{\Sigma}_t)}$ is injective. There is a certain subset of pairs (i, j) with $i \neq j \leq d$, and a homeomorphism $\gamma_{i,j} : a(\hat{\Sigma}_t^i) \rightarrow a(\hat{\Sigma}_t^j)$ which takes the image by a of the vertex of $\hat{\Sigma}_t^i$ to the image by a of the vertex of $\hat{\Sigma}_t^j$ such that two distinct elements $z_1, z_2 \in Z$ are in the same fibre of b if and only if $\gamma_{i,j}(z_1) = z_2$ for a certain pair (i, j) .

Using the mapping cylinder of the identifications a and b and Mayer-Vietoris sequences we deduce that the second homology group of $n_t(\hat{X}_t \cap \text{Int}\hat{H})$ coincides with the second homology group of $\hat{X}_t \cap \text{Int}\hat{H}$, which vanishes for being $\hat{X}_t \cap \text{Int}\hat{H}$ a disjoint union of cones. This proves that μ_0 is equal to 0, and hence that Γ is empty.

We only have to prove the claim. The claim is true for $t = 0$ since $\hat{H}_0$ is equal to $n_0^{-1}(B_{\epsilon/2})$. Let us prove it for any $t \in T'$. Define $R := X \cap (T' \times B_\epsilon) \setminus n(\text{Int}\hat{H})$. As the generic transversal Milnor number of f_t at any point of $\Sigma \cap R$ only depends on the connected component in which the point lies, using the technique of Section 2 of [3] we can find a neighbourhood V of $\Sigma \cap R$ in R such that $\partial_\pi V := \cup_{x \in \Sigma \cap R} \partial V_x$ is a manifold which meets transversely X_t for any t , and a trivialisation

$$\Psi_1 : V \rightarrow T' \times V_0$$

of $\tau|_V$. The restriction $\tau|_{R \setminus \dot{V}} : R \setminus \dot{V} \rightarrow T'$ is a proper topological submersion. Thus, taking T' contractible, it is a trivial fibration. Consider a trivialisation

$$\Psi_2 : R \setminus \dot{V} \rightarrow T' \times (R \setminus \dot{V})_0.$$

By Lemma 21 we can modify Ψ_2 so that it glues with Ψ_1 to get a trivialisation of $\tau|_R$. Then R_0 is homeomorphic to R_t and the claim for $t \neq 0$ follows from the claim for $t = 0$. $\square$

5.5. Existence of cuts for Σ . We prove that we can extend $n(\hat{H})$, being $\hat{H}$ the cut constructed in Section 5.3, to obtain a cut H for Σ such that $\text{Int}H \subset B(T, \eta)$. We extend $n(\hat{H})$ in two steps, first in a neighbourhood of Σ , and then out of it.

Step 1. Define

$$H_1 := E(n(\hat{H} \cap \hat{\Sigma}), \mu).$$

Because of the property (iv) of Definition 13 of the cut $\hat{H}$, we can ensure the equality

$$(5) \quad X \cap H_1 = n(\hat{H}) \cap E(\Sigma, \mu)$$

Step 2. We can consider a complex smooth line subbundle

$$\sigma : N \rightarrow X \setminus \Sigma$$

of the tangent bundle of $T \times \mathbb{C}^3$ that verifies

- it is a normal bundle to X ,
- the fibres of σ are contained in the fibres of τ and
- for any point $x \in E(\Sigma, \mu) \cap n(\hat{H}) \setminus \Sigma$ we have the inclusion $N_x \subset E_{\pi(x)}$.

Since the restriction of f to each fibre of σ is a local diffeomorphism at the origin (for being f non-singular at $X \setminus \Sigma$ and N a normal bundle), we can consider the metric obtained, in a neighbourhood of $X \setminus \Sigma$, as the pullback by f of a euclidean metric in $\mathbb{C}$. We can take a positive function $\nu : X \setminus \Sigma \rightarrow \mathbb{R}_+$ such that $N(X \setminus \Sigma, \nu)$ is embedded in $T \times \mathbb{C}^3$ as a tubular neighbourhood of $X \setminus \Sigma$ in $T \times \mathbb{C}^3 \setminus \Sigma$.

Consider the topological compact manifold with boundary $M := \overline{n(\hat{H}) \setminus E(\Sigma, \mu/2)}$. By compactness of M we can consider $\delta := \min\{\nu(x) : x \in M\}$. By the definition of the metric in N , the fibres $f_t^{-1}(s)$ are transverse to $N(M, \delta)$ for any $(t, s) \in T \times D_\delta$. Define

$$H_2 := N(M, \delta).$$

Thanks to Equality (5) and the third property of σ , the subspace $H_1 \cup H_2$ is a topological hypersurface of $T \times \mathbb{C}^3$. Define

$$H := H_1 \cup H_2.$$

If we have taken μ and δ small enough, we can get $H \subset B(T, \eta)$. Choose $\delta > 0$ such that $f_t^{-1}(s)$ is transverse to H (in the stratified sense when $s = 0$) for any $(t, s) \in T \times D_\delta$.

Property (iv) of Definition 12 follows directly from the properties of $\hat{H}$. Properties (i) – (ii) are satisfied by construction. Let us verify Property (iii).

We have to prove that $\text{Int}H(\delta)$ (defined as in Definition 14) is a bounded neighbourhood of T , the closure $\overline{\text{Int}H(\delta)}$ is a topological manifold and the restriction

$$\tau|_{\overline{\text{Int}H(\delta)}} : \overline{\text{Int}H(\delta)} \rightarrow T$$

a topological submersion. It is easy to see that it is enough to prove the statement locally over the base T . In the proof we will possibly have to shrink μ and δ . As T is compact such a shrinking only shall occur a finite number of times.

Take $t_0 \in T$ and $\epsilon > 0$, $\delta > 0$ such that S_ϵ is transverse to $f_{t_0}^{-1}(s)$ for any $s \in D_\delta$ and such that $B_\epsilon \cap H_{t_0} = \emptyset$. Take a neighbourhood T' of t_0 in T where these conditions are satisfied for any $t \in T'$.

Consider

$$K := f^{-1}(T' \times \partial D_\delta) \cup H.$$

It is clear that $\overline{\text{Int}_{T' \times \mathbb{C}^3} K} = \overline{\text{Int}H(\delta)}$.

Hence it is enough to construct a topological manifold L , included in K , which satisfies the following properties:

- The restriction $\tau|_L : L \rightarrow T'$ is a proper topological submersion whose fibre is a compact connected orientable manifold L_t ; thus, applying the Jordan-Brouwer Separation Theorem, for each $t \in T'$, the space $\mathbb{C}^3 \setminus L_t$ has two connected components. The bounded ones, that we call $\text{Int}L_t$, are neighbourhoods of the origin and $\partial \text{Int}L_t = L_t$.
- The intersection $\overline{\text{Int}L} \cap K$ coincides with L ; therefore $\overline{\text{Int}L} = \overline{\text{Int}K}$.

Let us construct L . Define

$$G_1 := (T' \times B_\epsilon) \cap f^{-1}(D_\delta),$$

$$L_1 := \partial G_1.$$

The restriction $\tau|_{L_1}$ is a proper topological submersion. Define

$$G_2 := G_1 \cup E(\text{Int}_\Sigma(H \cap \Sigma), \mu) \cap f^{-1}(D_\delta),$$

$$L_2 := \partial G_2.$$

Because of the transversality of Σ_t and $\mathbb{S}_\epsilon$ for every $t \in T'$, we can choose μ small enough (reducing δ accordingly) so that L_2 is a topological manifold and $\tau|_{L_2}$ is a topological submersion. Moreover $X \setminus \Sigma$ is transversal to L_2 .

Define

$$G := G_2 \cup N_{T'}(M, \delta),$$

$$L := \partial G.$$

As $X \setminus \Sigma$ is transversal to L_2 , if δ is small enough then L is a topological manifold. By construction it satisfies (a) and (b).

We have proved the following lemma

Lemma 25. *Let f_t be equisingular at the singular locus and equisingular at the normalisation. Given any positive function $\eta : T \rightarrow \mathbb{R}_+$, there exists a cut H of amplitude δ such that $\text{Int}H(\delta)$ is included in $B(T, \eta)$.*

5.6. Proof of Theorem 11. We restate and prove Theorem 11 in the following way,

Theorem 26. *Let f_t be equisingular at the singular locus and equisingular at the normalization. Consider a cut H for Σ of amplitude δ . There exists a trivialisation of τ relative to f*

$$\Theta : \text{Int}H(\delta) \rightarrow T \times (\text{Int}H)_0.$$

Proof. Applying Lemma 25 we construct a family of cuts H_i of amplitude δ_i (with $\delta_1 > \delta_2 > \dots$) such that

$$\text{Int}H(\delta) = \text{Int}H_1(\delta_1) \supset \dots \supset \text{Int}H_i(\delta_i) \supset \text{Int}H_{i+1}(\delta_{i+1}) \supset \dots$$

and

$$\bigcap_{i \in \mathbb{N}} \text{Int}H_i(\delta_i) = T.$$

Define

$$R_i := \overline{\text{Int}H_i(\delta_i) \setminus \text{Int}H_{i+1}(\delta_{i+1})}.$$

Following [3], let $\partial_\tau \text{Int}H_i(\delta_i)$ denote $\bigcup_{t \in T} \partial(\text{Int}H_i(\delta_i))_t$.

Lemma 27. *Any trivialisation*

$$\Xi_i : \partial_\tau \text{Int}H_i(\delta_i) \rightarrow T \times \partial(\text{Int}H_i(\delta_i))_0$$

of $\tau|_{\partial_\tau \text{Int}H_i(\delta_i)}$ relative to f can be extended to a trivialisation

$$\Theta_i : R_i \rightarrow T \times (R_i)_0$$

of $\tau|_{R_i}$ relative to f .

Proof of Lemma 27. We construct the trivialisation Θ_i piecewise. Split R_i in the following two spaces

$$P := \text{Int}H_i(\delta_{i+1}) \setminus \text{Int}H_{i+1}(\delta_{i+1})$$

$$Q := \text{Int}H_i(\delta_i) \setminus \text{Int}H_i(\delta_{i+1}).$$

Consider P as the union of two subsets, A and $B := \overline{P} \setminus A$ where A is defined below.

The function f restricted to any of the fibers of the complex vector bundle

$$\pi|_{E(\Sigma \cap R_i, \mu)} : E(\Sigma \cap R_i, \mu) \rightarrow \Sigma \cap R_i$$

is a holomorphic function with an isolated singularity at the origin whose Milnor number is independent of the considered fibre. By Section 2 of [3] there exist a cut S of amplitude δ_{i+1} (we would restrict δ_{i+1} if necessary) satisfying the inclusion

$$Y_{\text{int}}(S, \Sigma \cap R_i, \delta_{i+1}) \subset E(\Sigma \cap R_i, \mu).$$

We warn the reader that the notion of cut used in this paper is different from the one used in [3]; the cut S is a cut in the sense of [3] and following the notation of that paper, $Y_{\text{int}}(S, \Sigma \cap R_i, \delta_{i+1})$ stands for the union of the connected components of $E(\Sigma \cap R_i, \mu) \cap f^{-1}(D_{\delta_{i+1}}) \setminus S$ meeting $\Sigma \cap R_i$.

Define $A := Y_{\text{int}}(S, \Sigma \cap R_i, \delta_{i+1})$. By Theorem 11 of [3], there is a trivialisation

$$\Theta_A : A \rightarrow T \times A_0$$

of $\tau|_A$ relative to f . For being f_t equisingular at the critical locus,

$$F|_B : (B, A \cap B) \rightarrow T \times D_{\delta_{i+1}}$$

is a topological proper submersion of pairs; thus, we can extend the trivialisation of $F|_{A \cap B}$ given by Θ_A to a trivialisation of $F|_B$. This trivialisation induces a trivialisation of $\tau|_B$ relative to f

$$\Theta_B : B \rightarrow T \times B_0$$

whose restriction to $A \cap B$ coincides with Θ_A .

Define $\Theta_P : P \rightarrow T \times P_0$ glueing Θ_A and Θ_B . Analogously, the restriction

$$F|_Q : Q \rightarrow T \times (\overline{D_{\delta_i}} \setminus D_{\delta_{i+1}})$$

is a locally trivial fibration and there exist a homeomorphism

$$\beta : Q \rightarrow (Q \cap f^{-1}(\partial D_{\delta_i})) \times [0, 1]$$

which preserves the fibres of τ and carries fibres of f to fibres of $(f \circ p_1, p_2)$ being p_i the projections from $(Q \cap f^{-1}(\partial D_{\delta_i})) \times [0, 1]$ to the i th factor.

Consider the restrictions

$$\beta_0 : Q_0 \rightarrow Q_0 \cap f^{-1}(\partial D_{\delta_i}) \times [0, 1]$$

$$\alpha_0 : Q_0 \cap f^{-1}(\partial D_{\delta_{i+1}}) \rightarrow Q_0 \cap f^{-1}(\partial D_{\delta_i}) \times \{1\} \cong Q_0 \cap f^{-1}(\partial D_{\delta_i}).$$

Using the trivialisation Ξ_i of the frontier, we consider the composition

$$\gamma := (\Xi_i|_{Q \cap f^{-1}(\partial D_{\delta_i})}, Id_{[0,1]}) \circ \beta : Q \rightarrow T \times (Q_0 \cap f^{-1}(\partial D_{\delta_i})) \times [0, 1].$$

Composing with (Id_T, β_0^{-1}) we obtain a trivialisation of $\tau|_Q$ relative to f that extends $\Xi_i|_{f^{-1}(\partial D_{\delta_i}) \cap \text{Int}H_i(\delta_i)}$:

$$\tilde{\Xi}_i := (Id_T, \beta_0^{-1}) \circ \gamma : Q \rightarrow T \times Q_0$$

We will modify $\tilde{\Xi}_i$ in several steps, so that its restriction to $P \cap Q$ coincides with $\Theta_P|_{P \cap Q}$.

We extend first $\Theta_P|_{P \cap Q}$ to $Q \cap f^{-1}(\partial D_{\delta_{i+1}})$. For being $F|_{\text{Int}H_{i+1} \cap f^{-1}(\partial D_{\delta_{i+1}})}$ a proper topological submersion, there is a trivialisation of $\tau|_{\text{Int}H_{i+1} \cap f^{-1}(\partial D_{\delta_{i+1}})}$ relative to f that we call

$$\Upsilon : \text{Int}H_{i+1} \cap f^{-1}(\partial D_{\delta_{i+1}}) \rightarrow T \times (\text{Int}H_{i+1} \cap f^{-1}(\partial D_{\delta_{i+1}}))_0.$$

As it is clear that $(P \cap Q)_0 \cap H_{i+1}$ has a collar K in $(\text{Int}H_{i+1})_0 \cap f^{-1}(\partial D_{\delta_{i+1}})$ relative to f , applying Lemma 21 to Υ and $\Theta_P|_{P \cap Q}$, we obtain a trivialisation of $\tau|_{\text{Int}H_{i+1} \cap f^{-1}(\partial D_{\delta_{i+1}})}$ relative to f

$$\tilde{\Upsilon} : Q \cap f^{-1}(\partial D_{\delta_{i+1}}) \rightarrow T \times (Q_0 \cap f^{-1}(\partial D_{\delta_{i+1}})).$$

with $\tilde{\Upsilon}|_{P \cap Q} = \Theta_P|_{P \cap Q}$ and $\tilde{\Upsilon}|_{Q \cap f^{-1}(\partial D_{\delta_{i+1}}) \setminus \Upsilon^{-1}(K)} = \Upsilon|_{Q \cap f^{-1}(\partial D_{\delta_{i+1}}) \setminus \Upsilon^{-1}(K)}$.

In order that $\tilde{\Xi}_i$ can be glued with $\tilde{\Upsilon}$ it is enough that the self-homeomorphism of $T \times (Q_0 \cap f^{-1}(\partial D_{\delta_{i+1}}))$, defined by

$$h := \tilde{\Xi}_i|_{Q \cap f^{-1}(\partial D_{\delta_{i+1}})} \circ \tilde{\Upsilon}^{-1}$$

is the identity.

By construction we have that h_0 is the identity. By Lemma 15 there is an isotopy

$$\phi : T \times (Q_0 \cap f^{-1}(\partial D_{\delta_{i+1}}) \times [0, 1]) \rightarrow T \times (Q_0 \cap f^{-1}(\partial D_{\delta_{i+1}}) \times [0, 1])$$

such that $\phi_0 = Id$ and $\phi_1 = h^{-1}$. Substituting γ by

$$(Id_T, \alpha_0^{-1}, Id_{[0,1]}) \circ \phi \circ (Id_T, \alpha_0, Id_{[0,1]}) \circ \gamma$$

in the definition of $\tilde{\Xi}_i$ we obtain that $\tilde{\Xi}$ extends $\tilde{\Upsilon}$.

Now, we can define Θ glueing Θ_P and $\tilde{\Xi}_i$. As there is a collar of $H_i(\delta_i)_0$ in $(R_i)_0$ relative to f and to $f^{-1}(\partial D_{\delta_i}) \cap (R_i)_0$, we can apply Lemma 21 to modify Θ so that it extends Ξ_i . □

We construct Θ_1 following the proof of the previous Lemma, obtaining Θ_Q directly from the locally trivial fibration $(\tau, f) : Q \rightarrow T \times (D_{\delta_i} \setminus D_{\delta_{i+1}})$ where T is contractible.

Using the lemma we construct inductively a family of trivialisations

$$\Theta_i : R_i \rightarrow T \times (R_i)_0$$

of $\tau|_{R_i}$ relative to f which glue to a homeomorphism

$$\Theta^* : \text{Int}H(\delta) \setminus T \rightarrow T \times (\text{Int}H(\delta) \setminus T)_0.$$

This homeomorphism can be extended to the desired trivialisation simply defining $\Theta|_T := (\tau, O)$. □

6. EQUISINGULARITY OF PARAMETRISED SURFACES

Using the theory developed above, we are going to study topological $\mathcal{A}$ -equisingularity of holomorphic families of surface parametrisations. By a parametrisation we understand a finite, generically one to one mapping from $\mathbb{C}^2$ to its image in $\mathbb{C}^3$. A parametrisation necessarily coincides with the normalisations of its image.

Let T be a contractible and compact complex manifold. A family of parametrisations over T is a finite holomorphic germ

$$\varphi : T \times (\mathbb{C}^2, O) \rightarrow T \times (\mathbb{C}^3, O),$$

such that $\varphi_t(x) := \varphi(t, x)$ is a parametrisation.

Definition 28. *The family φ_t is $\mathcal{A}$ -topologically equisingular if there are homeomorphism germs*

$$\begin{aligned} \beta : T \times (\mathbb{C}^2, O) &\rightarrow T \times (\mathbb{C}^2, O) \\ \gamma : T \times (\mathbb{C}^3, O) &\rightarrow T \times (\mathbb{C}^3, O) \end{aligned}$$

which preserve the fibres of the respective projections to T and such that we have the equality

$$\gamma \circ \varphi \circ \beta = (Id_T, \varphi_0).$$

If in addition the restrictions $\beta|_{T \times \mathbb{C}^2 \setminus T}$, $\gamma|_{T \times \mathbb{C}^3 \setminus \Sigma}$ and $\gamma|_{\Sigma \setminus T}$ are smooth, we say that the family φ_t is $\mathcal{A}$ -piecewise smoothly equisingular.

Lemma 29. *Let $X \subset T \times \mathbb{C}^3$ be a family of surfaces and $n : \hat{X} \rightarrow X$ be the normalisation. If $\hat{X}_t = \mathbb{C}^2$ for any $t \in T$, and X_t is equisingular at the singular locus, then the family $\hat{\Sigma}_t$ is a reduced μ -constant family of plane curves.*

Proof. By the second condition of equisingularity at the singular locus the space $X \setminus T$ is equisingular at Σ . Thus the restriction $n : T \times (\mathbb{C}^2 \setminus \{O\}) \rightarrow X \setminus T$ is the simultaneous resolution of the singularities of $X \setminus T$. This implies that the restriction

$$n : \hat{\Sigma} \setminus \hat{T} \rightarrow \Sigma \setminus T$$

is an unramified covering. Hence, the first condition of equisingularity at the singular locus, implies that $\hat{\Sigma} \setminus \hat{T}$ is a topological trivial fibration over T .

The conclusion of the Lemma is now easy to obtain. $\square$

Theorem 30. *Let φ_t be a family of parametrisations. If the family is topologically $\mathcal{A}$ -equisingular, then the Milnor number of $\hat{\Sigma}_t$ at the origin is constant. If $L\hat{e}$'s Conjecture is true the converse holds.*

If for a single $t \in T$ each of the generic transversal singularities of $\varphi_t(\mathbb{C}^2)$ has at least one smooth branch, then the converse holds without $L\hat{e}$'s Conjecture.

Proof. Let $f : T \times \mathbb{C}^3 \rightarrow \mathbb{C}$ be a reduced equation of the image of φ . The topological $\mathcal{A}$ -equisingularity of φ_t implies the embedded topological equisingularity of f_t . This implies that f_t is abstractly equisingular in the strong sense. By Proposition 6 and Lemma 3 (vii), we have that the Milnor number of $\hat{\Sigma}_t$ is constant.

Assuming the constancy of the Milnor number of $\hat{\Sigma}_t$ and the fact that $\hat{X}_t$ is $\mathbb{C}^2$ for any $t \in T$, we obtain that f_t is equisingular at the normalisation. By Proposition 9, the family f_t is equisingular at the singular locus. By Theorem 26, there is a family of homomorphisms $\gamma_t : (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}^3, 0)$ such that $\gamma_t(X_0) = X_t$. Using Lemma 5, we can construct $\beta_t : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$ such that

$$\gamma_t \circ \varphi_t \circ \beta_t = \varphi_0.$$

$\square$

Corollary 31. *Let φ_t be a family of parametrisations. If the generic transversal singularities of $\varphi_t(\mathbb{C}^2)$ are of Morse type, then the family is topologically $\mathcal{A}$ -equisingular if and only if the Milnor Number of $\hat{\Sigma}_t$ is independent of $t \in T$.*

Corollary 32 (Calleja-Bedregal, Houston, Ruas). *Let φ_t be a family of finitely determined mapping germs. The family is topologically $\mathcal{A}$ -equisingular if and only if the Milnor Number of $\hat{\Sigma}_t$ is independent of $t \in T$.*

Proof. In the finitely determined case, mapping germs are parametrisations and transversal singularities are of Morse type. $\square$

Theorem 33. *Let φ_t a family of parametrisations and f_t the reduced equations of the X_t . The constancy of the generic Lê numbers of f_t implies topological $\mathcal{A}$ -equisingularity of φ_t .*

Proof. The constancy of Lê numbers implies equisingularity at the critical locus as it was proved in [3] Theorem 48. By Lemma 29 the family $\hat{\Sigma}_t$ is μ -constant and hence φ_t is equisingular at the normalisation. Once we have equisingularity at the singular locus and equisingularity at the normalisation, we reproduce the proof of Theorem 30. $\square$

7. THE PIECEWISE SMOOTH EQUISINGULARITY

The technique of this article can be readily adapted to get the analogous results for the piecewise smooth case.

Theorem 34. *Let f_t be equisingular at the singular locus and piecewise smooth equisingular at the normalisation, then it is piecewise smooth R -equisingular.*

Corollary 35. *Let φ_t a family of parametrisations. If Lê's Conjecture is true and the Milnor number of $\hat{\Sigma}_t$ at the origin is constant, then the family is piecewise smoothly $\mathcal{A}$ -equisingular.*

If for a single $t \in T$ each of the generic transversal singularities of $\varphi_t(\mathbb{C}^2)$ has at least a smooth branch, then the conclusion holds without Lê's Conjecture.

Proof. Since $\hat{X}_t = \mathbb{C}^2$ for any $t \in T$, the constancy of Milnor number of $\hat{\Sigma}_t$, implies piecewise smooth equisingularity at the normalisation. $\square$

Observe that Lê's Conjecture implies that the notions of topological $\mathcal{A}$ -equisingularity and piecewise smooth $\mathcal{A}$ -equisingularity are equivalent since topological $\mathcal{A}$ -equisingularity implies the constancy of the Milnor number of $\hat{\Sigma}_t$.

Corollary 36. *Let φ_t a family of parametrisations. If the generic transversal singularities of $\varphi_t(\mathbb{C}^2)$ are of Morse type, then the family is piecewise smoothly $\mathcal{A}$ -equisingular if and only if the Milnor Number of $\hat{\Sigma}_t$ is independent of $t \in T$.*

Corollary 37. *Let $\varphi_t : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^3, 0)$ be a holomorphic family of finitely determined mapping germs. The family is piecewise smoothly $\mathcal{A}$ -equisingular if and only if the Milnor Number of $\hat{\Sigma}_t$ is independent of $t \in T$.*

REFERENCES

- [1] R.O. Buchweitz, G.M. Greuel. *The Milnor number and deformations of complex curve singularities.* Invent. Math. **58** no. 3, (1980), 241-281.
- [2] Calleja-Bedregal, Houston, Ruas. *Topological triviality of families of singular surfaces.* Preprint (2006).
- [3] J. Fernández de Bobadilla. *Topological equisingularity of function germs with 1-dimensional critical locus.* ArXiv: MathAG/0603508.
- [4] J. Fernández de Bobadilla. *A reformulation of Lê's Conjecture.* To appear in Indagationes Math.

- [5] T. Gaffney. *Polar multiplicities and equisingularity of map germs*. Topology **32** no. 1, (1993), 185-223.
- [6] T. de Jong, T., D. van Straten. *D. Deformations of the normalization of hypersurfaces*. Math. Ann. **288** (1990), no. 3, 527-547.
- [7] T. de Jong, T., D. van Straten. *A deformation theory for nonisolated singularities*. Abh. Math. Sem. Univ. Hamburg **60** (1990), 177-208.
- [8] T. de Jong, T., D. van Straten. *On the base space of a semi-universal deformation of rational quadruple points*. Annals of Math. **134**, (1991), 653-678.
- [9] R. Kirby, L.C. Siebenmann. *Foundational essays on topological manifolds, smoothings and triangulations*. Annals of Mathematical Studies **88** (1977).
- [10] *Noeuds, Tresses et Singularités*. Proceedings of the seminar held in Plans-sur-Bex, March 1982. C. Weber (ed). Monographies de L'Enseignement Mathématique **31**, (1983).
- [11] D. Massey. *The Lê varieties I*. Invent. Math. **99** (1990), 357-376.
- [12] D. Massey. *The Lê varieties II*. Invent. Math. **104** (1991), 113-148.
- [13] D. Massey. *Lê cycles and hypersurface singularities*. Lecture Notes in Mathematics **1615**. Springer (1995).
- [14] D. Mond. *Some remarks on the geometry and classification of germs of maps from surfaces to 3-space*. Topology **26** no. 3, (1987), 361-383.
- [15] D. Mumford *The topology of normal singularities of an algebraic surface and a criterion for simplicity*. Inst. Hautes udes Sci. Publ. Math. **9**, (1961), 5-22.

DEPARTAMENTO DE ÁLGEBRA. FACULTAD DE CIENCIAS MATEMÁTICAS. UNIVERSIDAD COMPLUTENSE. PLAZA DE CEINCIAS 3. 28040 MADRID. SPAIN.

E-mail address: javier@mat.uned.es

DEPARTAMENTO DE ÁLGEBRA. FACULTAD DE CIENCIAS MATEMÁTICAS. UNIVERSIDAD COMPLUTENSE. PLAZA DE CEINCIAS 3. 28040 MADRID. SPAIN.

E-mail address: maria.pe@mat.ucm.es